

In: Environmental Pollution: New Research
Editor: Rachel H. Plattenberg, pp. 121-139

ISBN: 1-60021-285-9
© 2007 Nova Science Publishers, Inc.

Chapter 4

MANAGEMENT OF IRRIGATED AGRICULTURE TO INCREASE ORGANIC CARBON STORAGE IN SOILS

James A. Entry^{1*}, R.E. Sojka^{1**} and Glenn E. Shewmaker^{2}***

¹ USDA Agricultural Research Service, Northwest Irrigation and Soils Research
Laboratory, 3793 North, 3600 East, Kimberly, Idaho 83341

³ University of Idaho, Research and Extension Center, Twin Falls, Idaho 83303-1827.

Abstract

Increasing the amount of C in soils may be one method to reduce the concentration of CO₂ in the atmosphere. We measured organic C stored in southern Idaho soils having long term cropping histories that supported native sagebrush vegetation (NSB), irrigated moldboard plowed crops (IMP), irrigated conservation -chisel- tilled crops (ICT) and irrigated pasture systems (IP). The CO₂ emitted as a result of fertilizer production, farm operations and CO₂ lost via dissolved carbonate in irrigation water, over a 30 year period, was included. Net organic C in ecosystems decreased in the order IP>ICT>NSB>IMP. In this study, if NSB were converted to IMP, 0.15 g C m⁻² would be emitted to the atmosphere, but if converted to IP 3.56 g C m⁻² could be sequestered. If IMP land were converted to ICT, 0.95 g C m⁻² could be sequestered in soil and if converted to IP 3.71 g C m⁻² could be sequestered. There are 2.6x10⁸ ha of land worldwide presently irrigated. If irrigated agriculture were expanded 10% and the same amount of rainfed land were converted back to native grassland, an increase of 3.4x10⁹ Mg C (5.9% of the total C emitted in the next 30 yr) could potentially be sequestered. The total projected release of CO₂ is 5.7x10¹⁰ Mg C worldwide during the next 30 years. Converting rainfed agriculture back to native vegetation while modestly increasing areas in irrigated agriculture could have a significant impact on CO₂ atmospheric concentrations while maintaining or increasing food production.

* E-mail address: jentry@kimberly.ars.pn.usbr.gov; Phone (208) 423 - 6553, Fax (208) 423-6555. Corresponding author.

** E-mail address: sojka@kimberly.ars.pn.usbr.gov; Phone (208) 423 - 6562, Fax (208) 423-6555

*** E-mail address: gshew@uidaho.edu; Phone (208) 736-3600, Fax (208) 736-0843

Abbreviations

C	carbon;
NSB	native sagebrush;
IMP	irrigated moldboard plowed crop land;
ICT	crop land managed with conservation (chisel) tillage;
IP	irrigated pasture;
C _S	carbon projected to sequestered in soils worldwide over the next 30 years;
C _{EW}	carbon projected to be emitted worldwide over the next 30 years.

Introduction

In 1992, nearly all countries of the world signed the Framework Convention on Climate change. Its long term goal is to stabilize the concentration of greenhouse gases in the atmosphere at concentrations that should prevent dangerous anthropogenic interference with the climate system. To stabilize or reduce CO₂ concentrations, the gas must be transferred from the atmosphere to marine or terrestrial ecosystems. Processes or activities that remove greenhouse gases from the atmosphere are defined as sinks in the Framework Convention. In the 1997 Kyoto Protocol, agricultural soils are specifically recognized in the list of potential sinks of greenhouse gases.

Soils are the largest pool of C in the terrestrial environment (Jobbagy and Jackson, 2000; Schlesinger, 1990; 1995). The amount of C stored in soils is twice the amount of C in the atmosphere and three times the amount of C stored in living plants (Schlesinger, 1990; 1995; Kimble and Stewart, 1995), therefore, a change in the size of the soil C pool could significantly alter the atmospheric CO₂ concentration (Wang et al., 1999). The current concentration of C in soils reflects the balance between past C accumulation and loss. Accumulation of C in soils is derived from litter and root input, while losses result from microbial degradation of organic matter, eluviation and erosion (Entry and Emmingham, 1998). At equilibrium, the rate and amount of C added to the soil via vegetation are equal to the rate and amount of C lost through organic matter degradation and other losses (Henderson, 1995). Within limits, soil C increases with increasing soil water and decreasing temperature (Hontoria et al., 1999; Wang et al., 1999; Burke et al., 1989). The effect of soil water is much greater than the effect of soil temperature (Birch and Friend, 1956; Hontoria et al., 1999; Liski et al., 1999). Increasing water within temperature zones can increase plant production and, thus, C input to soils via increased plant litter and root production (Liski et al., 1999).

Land use changes can impact the amount of C stored in the soil by altering C inputs and losses. In forest, grassland and wetland ecosystems, conversion of native vegetation to agricultural cropping has resulted in substantial C transfer to the atmosphere as a result of loss of climax vegetation to the lower equilibrium C concentration in soil (Lal et al., 1999; Wang et al., 1999; Cambardella and Elliot, 1992; Johnson, 1992). In arid and semi-arid environments plant survival and growth is limited by available water and irrigation is required to increase plant production to the point where crops become economically viable. Irrigation also increases C input to soils via increased litter and root production.

When assessi
soils, one needs to
manufacture, stor
irrigation water, C
carbonate in irri
(1999) used a f
fertilization rate i
yr⁻¹ is approxima
realistic rate is 10
from pumping ir
using gasoline, to
addition, C may
semiarid regions
soil, CaCO₃ can p
0.05 g L⁻³ dissolv
in plant C is 2000
CO₂ released via :

Farm manag
reduced the amou
(West and Marlar
1991). Intensively
improved grazin
(Follett, 2001; Br
semiarid lands by
storage in soils. T
moldboard plowe
sequester addition
management polic
have estimated.

Materials an

Site Descriptio

The study area is
and 114° 20' 00"
from 860 m to 13
1998). The clima
precipitation rang
through March (C
typically well-dra
Vegetation throug
(*Artemisia trideni*
var. *wyomingensi*.
secunda J. Presl

When assessing the potential of irrigation of arid or semiarid land to increase C storage in soils, one needs to assess C loss from CO₂ emitted to the atmosphere as a result of 1) fertilizer manufacture, storage, transport and application, 2) fossil-fuel CO₂ emitted from pumping irrigation water, 3) farm operations, such as tillage and planting, and 4) CO₂ lost via dissolved carbonate in irrigation water (West and Marland, 2001; Schlesinger, 1999). Schlesinger (1999) used a fertilization value of 336 kg N ha⁻¹ yr⁻¹, which is an unusually high fertilization rate in U.S. farms. The CO₂ released during fertilizer production of 336 kg N ha⁻¹ yr⁻¹ is approximately 16.7 g C m⁻² yr⁻¹ (Schlesinger, 1999). It has been noted that a more realistic rate is 100-150 kg N ha⁻¹ yr⁻¹ (West and Marland, 2001). Carbon dioxide released from pumping irrigation water in the United States ranges from 126 kg C ha⁻¹ yr⁻¹, when using gasoline, to 266 kg C ha⁻¹ yr⁻¹ when using electricity (West and Marland, 2001). In addition, C may be lost as CO₂ from the irrigation water itself. Irrigation water in arid and semiarid regions often contains as much as 1% dissolved CO₂. When water is applied to the soil, CaCO₃ can precipitate, releasing CO₂ into the atmosphere. If irrigation water containing 0.05 g L⁻³ dissolved Ca is used to irrigate crops in semi-arid climates, the calculated increase in plant C is 2000 g C m⁻² per year over C contained in native soils and vegetation. The net CO₂ released via irrigation water is calculated to be 8.4 g C m⁻² yr⁻¹ (Schlesinger, 1999).

Farm management practices, including conservation tillage and erosion control, have reduced the amount of CO₂ emitted to the atmosphere in both Canada and the United States (West and Marland, 2001; Janzen et al., 1997; Paustian et al., 1997; Rasmussen and Collins, 1991). Intensively managed crop or pasture lands has potential for C gain through the use of improved grazing regimes, improved fertilization practices and irrigation management (Follett, 2001; Bruce et al., 1999). We hypothesize that increasing plant growth on arid and semiarid lands by conversion to irrigated agriculture is one method that may increase C storage in soils. The objective of this research was to determine if land managed as irrigated moldboard plowed crops converted to irrigated conservation tillage or irrigated pasture could sequester additional C. We use our findings to pose several possible scenarios of altered land management policies that could favor global C sequestration based on the C budgets that we have estimated.

Materials and Methods

Site Descriptions

The study area is located on the Snake River Plain, between 42° 30' 00" and 43° 30' 00" N. and 114° 20' 00" and 116° 30' 00" W. The sites occur across an elevational gradient ranging from 860 m to 1300 m. The area is classified as a temperate semi-desert ecosystem (Bailey, 1998). The climate is typified by cool, moist winters and hot, dry summers with annual precipitation ranging from 175 to 305 mm, two-thirds of which occurs during October through March (Collett, 1982). Average annual temperature ranges from 9 to 10 °C. Soils are typically well-drained loams and silt loams derived from loess deposits overlying basalt. Vegetation throughout the general area was historically dominated by basin big sagebrush (*Artemisia tridentata* var. *tridentata* Nutt.), Wyoming big sagebrush (*Artemisia tridentata* var. *wyomingensis* Nutt.), and perennial bunch grasses, including Sandberg bluegrass (*Poa secunda* J. Presl), bottlebrush squirreltail (*Elymus elymoides* Raf. Swezy.), bluebunch

wheatgrass (*Pseudoroegneria spicata* Pursh. A. Love), and Thurber's needlegrass (*Achnatherum thurberianum* (Piper) Barkworth).

Experimental Design

The experiment was arranged in a completely randomized design (Kirk, 1982). Soil samples were taken from: 1) three sites supporting native sagebrush vegetation (NSB) located near agricultural land in Southern Idaho (each site supported a basin big sage and a Wyoming big sage vegetation type); 2) three sites that were formerly crop land and converted to and maintained as irrigated pasture (IP) for the past 30 years; 3) three sites that were irrigated crop land and have been managed with conservation tillage (ICT) for the past 8 years; and 4) three irrigated agricultural crop lands in moldboard plowing systems (IMP) that were each growing a) alfalfa, b) wheat, c) potato and d) beans. There were four treatments (NSB, IMP, ICT, and IP) x three sites for each treatment x five cores taken within each treatment at each site (replications) x four soil depths (0-5 cm, 5-15 cm, 15-30 and 30-100 cm). We took a total of 240 samples.

Native Vegetation Sagebrush Sites

Native sagebrush sites were vegetated with native steppe vegetation and a low composition of exotic annual grasses. Sites were chosen for this study based on a history of no livestock grazing (BLM, Bruneau Resource Area, unpublished data). All study sites had 5-10 % slope and were on areas that supported basin big sagebrush or Wyoming big sagebrush or communities (Table 1). Soil was classified as a fine, montmorillonitic, mesic Xerollic Haplargid on the Brown's Creek site, a coarse-loamy, mixed non-acid, mesic Xeric Torriorthents on the Simco site and a loamy, mixed, mesic lithic Xerollic Camborthids on the Kuna Butte site (Collett, 1982).

Irrigated Pasture Sites

Three irrigated pastures were selected that were formerly crop land and converted to and maintained as irrigated pasture (IP) for the past 30 years. The Buhl site was vegetated with Kentucky bluegrass (*Poa pratensis* L.)-orchardgrass (*Dactylis glomerata* L.) on a Rakane-Blacknest soil complex, fine-loamy, mixed, mesic Xerollic Durargids soil. The Gooding site was vegetated with smooth brome (*Bromus inermis* Leys.)-orchardgrass on a Paulville-Idow soil complex, fine-loamy, mixed, mesic Xerollic Haplargid soil. The Kimberly site was vegetated with smooth brome-orchardgrass pasture on a Portneuf soil, coarse-silty, mixed, superactive, mesic Durinodic Xeric Haplocalcid soil. Grazing rates on these pastures were 10-12 animal unit months yr⁻¹.

Table 1. Location and characteristics of study sites.

Site Name	Latitude Longitude (10' block)	Treatment	Elevation (m)	Site Description	Last Known Fire(s)	Current Dominant Vegetation
Bowns Creek	43° 20' 00" 115° 56' 00"	Native	1010	Artemisia tridentata Basin / Pseudoroegneria (Pursh) A. Love. Artemisia tridentata Wyoming / Pseudoroegneria	unknown (>50 years)	Artemisia tridentata Basin, Wyoming / Pseudoroegneria

er's needlegrass

982). Soil samples (NSB) located near and a Wyoming big converted to and were irrigated crop years; and 4) three were each growing SB, IMP, ICT, and treatment at each site. We took a total of

ow composition of ry of no livestock had 5-10 % slope big sagebrush or ic, mesic Xerollic acid, mesic Xeric Camborthids on the

i converted to and was vegetated with : L.) on a Rakane-. The Gooding site on a Paulville-Idow Kimberly site was coarse-silty, mixed, e pastures were 10-

Table 1. Location and characteristics of study sites.

Site Name	Latitude Longitude (10' blocks)	Treatment	Elevation (m)	Site Description	Last Known Fire(s)	Current Dominant Vegetation
Bowns Creek	43° 20' 00" 115° 56' 00"	Native	1010	<i>Artemisia tridentata</i> Basin / <i>Pseudoroegneria</i> (Pursh) A. Love. <i>Artemisia tridentata</i> Wyoming / <i>Pseudoroegneria</i> Soil Type: Xeric Haplargid	unknown (>50 years)	<i>Artemisia tridentata</i> Basin, Wyoming / <i>Pseudoroegneria</i> <i>spicata</i> / <i>Poa secunda</i>
Simco	43° 22' 00" 115° 51' 00"	Native	1095	<i>Artemisia tridentata</i> Basin / <i>Pseudoroegneria</i> <i>Artemisia tridentata</i> Wyoming / <i>Pseudoroegneria</i> Soil Type: Xeric Torriorthent	unknown (>50 years)	<i>Artemisia tridentata</i> Basin, Wyoming / <i>Pseudoroegneria</i> <i>spicata</i> / <i>Poa secunda</i>
Kuna West	43° 27' 00" 116° 27' 00"	Native	950	<i>Artemisia tridentata</i> Basin / <i>Pseudoroegneria</i> <i>Artemisia tridentata</i> Wyoming / <i>Pseudoroegneria</i> Soil Type: Xerollic Camborthid	unknown (>50 years)	<i>Artemisia tridentata</i> Basin, Wyoming / <i>Poa secunda</i>
Kimberly Pasture	42° 00' 00" 114° 30' 00"	Irrigated Pasture	1210	<i>Dactylis glomerata</i> L. / <i>Bromus inermis</i> Lyess. Soil Type: Xeric Haplocalcid	(>100 years)	50 % <i>Dactylis glomerata</i> . 50 % <i>Bromus inermis</i>
Gooding Pasture	42° 46' 00" 114° 37' 00"	Irrigated Pasture	1070	<i>Bromus inermis</i> / <i>Dactylis glomerata</i> Soil Type: Xerollic Haploargid	(>100 years)	50 % <i>Dactylis glomerata</i> 50 % <i>Bromus inermis</i>
Buhl Pasture	42° 30' 00" 114° 30' 00"	Irrigated Pasture	1073	<i>Poa pratensis</i> L. / <i>Dactylis glomerata</i> . Soil Type: Xerollic Durargids	(>100 years)	50 % <i>Poa pratensis</i> 50 % <i>Dactylis glomerata</i>
Kimberly	42° 00' 00" 114° 30' 00"	Irrigated moldboard plow	1260	Rotations of: <i>Phaseolus vulgaris</i> L. / <i>Triticum aestivum</i> L. / <i>Solanum tuberosum</i> L. / <i>Medicago sativa</i> L. Durinodic Xeric Haplocalberid	(>100 years)	<i>Phaseolus vulgaris</i> L. / <i>Triticum</i> <i>aestivum</i> L. / <i>Solanum tuberosum</i> L. / <i>Medicago</i> <i>sativa</i> L.
University of Idaho	42° 46' 00" 114° 37' 00"	Irrigated moldboard plow	1220	Rotations of: <i>Phaseolus vulgaris</i> L. / <i>Triticum aestivum</i> L. / <i>Solanum tuberosum</i> L. / <i>Medicago sativa</i> L. Durinodic Xeric Haplocalcid	(>100 years)	<i>Phaseolus vulgaris</i> L. / <i>Triticum</i> <i>aestivum</i> L. / <i>Solanum tuberosum</i> L. / <i>Medicago</i> <i>sativa</i> L.
South Farm	42° 30' 00" 114° 30' 00"	Irrigated moldboard plow	1200	Rotations of: <i>Phaseolus vulgaris</i> L. / <i>Triticum aestivum</i> L. / <i>Solanum tuberosum</i> L. / <i>Medicago sativa</i> L. Durinodic Xeric Haplocalcid	(>100 years)	<i>Phaseolus vulgaris</i> L. / <i>Triticum</i> <i>aestivum</i> L. / <i>Solanum tuberosum</i> L. / <i>Medicago</i> <i>sativa</i> L.

Table 1. Continued

Site Name	Latitude Longitude (10' blocks)	Treatment	Elevation (m)	Site Description	Last Known Fire(s)	Current Dominant Vegetation
Kimberly	42° 00' 00" 114° 30' 00"	Irrigated conservation tillage	1260	Rotations of: <i>Phaseolus vulgaris</i> L. / <i>Triticum aestivum</i> L. / <i>Solanum tuberosum</i> L. / <i>Medicago sativa</i> L. / Durinodic Xeric Haplocamberid	(>100 years)	<i>Phaseolus vulgaris</i> L. / <i>Triticum aestivum</i> L. / <i>Solanum tuberosum</i> L. / <i>Medicago sativa</i> L.
University of Idaho	42° 46' 00" 114° 37' 00"	Irrigated conservation tillage	1220	Rotations of: <i>Phaseolus vulgaris</i> L. / <i>Triticum aestivum</i> L. / <i>Solanum tuberosum</i> L. / <i>Medicago sativa</i> L. / Durinodic Xeric Haplocalcid	(>100 years)	<i>Phaseolus vulgaris</i> L. / <i>Triticum aestivum</i> L. / <i>Solanum tuberosum</i> L. / <i>Medicago sativa</i> L.
South Farm	42° 30' 00" 114° 30' 00"	Irrigated conservation tillage	1200	Rotations of: <i>Phaseolus vulgaris</i> L. / <i>Triticum aestivum</i> L. / <i>Solanum tuberosum</i> L. / <i>Medicago sativa</i> L. / Durinodic Xeric Haplocalcid	(>100 years)	<i>Phaseolus vulgaris</i> L. / <i>Triticum aestivum</i> L. / <i>Solanum tuberosum</i> L. / <i>Medicago sativa</i> L.

Irrigated Con.

Three sites with 1 L.), potato (*Solar* were located on Irrigation and So Center. Soil on a Xeric Haplocalc approximately 13 3.0% (Table 1).

Sampling Proc

Soil cores were (August), autumn spring, summer a irrigation, tillage field. Separate 10 cm, 5-15 cm, 1. measured separat amount of materi 1998).

Carbon in Soil

Concentration of Black procedure of the 0-100 cm correction for soi depth, divided int Bulk density was hours by the volu collected and sep dried at 80 °C for vegetation was de in the above grou free basis (Nelso Western United S

Calculations

Concentration of bulk density meas was converted to

Irrigated Conservation Tillage and Crop Sites

Three sites with fields rotating among alfalfa (*Medicago sativa* L.), wheat (*Triticum aestivum* L.), potato (*Solanum tuberosum* L.) and beans (*Phaseolus vulgaris* L.) were sampled. All sites were located on fields managed by USDA Agricultural Research Service's Northwest Irrigation and Soils Research Laboratory or the University of Idaho, Research and Extension Center. Soil on all sites was classified as a coarse-silty, mixed, superactive, mesic Durinodic Xeric Haplocalcid, with 0.1-0.21 g/g clay and 0.6-0.75 g/g silt, and organic matter of approximately 13 g kg⁻¹. The soil has a pH of 7.6-8.0. Slope on these sites ranges from 1.0 to 3.0% (Table 1).

Sampling Procedures

Soil cores were taken from each site during winter (January), spring (April), summer (August), autumn (November) in 1999. We sampled the top 1 m of soil each season (winter, spring, summer and autumn) to determine if the amount of C in soil would be affected by irrigation, tillage and vegetation. Sampling locations were randomly chosen at each site or field. Separate 10 cm diameter replicate cores were randomly taken and partitioned into 0-5 cm, 5-15 cm, 15-30 and 30-100 cm depths. Roots greater than 1.0 cm diameter were measured separately. Carbon in above-ground vegetation was estimated by measuring the amount of material in 10 separate 1.0 m² areas in each site or field (Entry and Emmingham, 1998).

Carbon in Soil and Above-ground Vegetation

Concentration of organic C in each sample of mineral soil was determined by the Walkley-Black procedure and loss on ignition (Nelson and Sommers, 1996). The amount of C per ha⁻¹ of the 0-100 cm of mineral soil was calculated assuming 0.44 g C g⁻¹ organic matter with correction for soil bulk density. Ten separate 10 cm diameter soil cores were taken to a 1.0 m depth, divided into 0-5 cm, 5-15 cm, 15-30 and 30-100 cm depths to determine bulk density. Bulk density was measured by dividing by the oven dry weight after drying at 105 °C for 48 hours by the volume of the sample (Blake and Hartage, 1982). Above-ground vegetation was collected and separated into sage, grass, forbs, herbs and duff. Above-ground material was dried at 80 °C for 48 h, weighed and ground to pass a 1 mm opening. Carbon in above-ground vegetation was determined by loss on ignition (Nelson and Sommers, 1996). The amount of C in the above ground material was assumed to contain 0.44 g C g⁻¹ organic matter on an ash free basis (Nelson and Sommers, 1996). Calculations estimating C stored in soils in the Western United States and worldwide are based on the Walkley Black procedure.

Calculations

Concentration of organic C determined by the Walkley-Black procedure was converted, using bulk density measurements, to a meter square basis to a depth of 1 meter. Organic C in kg m⁻² was converted to Mg C ha⁻¹ by multiplying by 10,000 (land area) and dividing by 1,000 (C

weight), which is a 1: 10 conversion. We divided the resulting number by 10 to account for a 10 % conversion of one treatment (land area). The amount of C sequestered in the Pacific Northwestern United States, the 11 western states in the United States and worldwide was estimated by multiplying Mg C ha⁻¹ by the number of hectares of irrigated land in each area. There are 9,055, 979 ha of land in irrigated crop land in the Pacific Northwest, 24,322,029 ha in the Western U.S. and 260, 000,000 ha worldwide (Bucks et al., 1990; Tribe, 1994; Howell, 2000). The carbon sequestered worldwide (C_S) relative to the amount of C projected to be emitted worldwide during the next 30 yr (C_{EW}) was calculated by dividing the Mg C sequestered in each treatment x area by the total projected worldwide release of CO₂ - C during the next 30 years (5.7 x 10¹⁰ Mg C) multiplied by 100.

Statistical Analysis

All data were subjected to a one way vegetation type analysis of variance (ANOVA) for a completely randomized design (Snedecor and Cochran, 1980; Kirk, 1982). Residuals were normally distributed with constant variance. Statistical Analysis Software programs (SAS Institute Inc. 1996) were used to conduct the analysis of variance. Significance of treatment means were determined at P < 0.05 with the Least Square Means test.

Results and Discussion

Site Specific Findings

Statistical comparisons in the ANOVA showed that soil bulk density, soil C, site C, net C in soil, net site C in site x vegetation interactions are not significant at P<0.05. Therefore, results are discussed with respect to vegetation differences (Snedecor and Cochran 1980; Kirk 1982). Bulk density was greater in soils in IP, ICT and IMP than in NSB soils. Bulk density was less in the NSB 0-5 cm soil depth than the 5-15, 15-30 and 30-100 cm depths and all other soils (Table 2). Soil C was greater in the NSB 0-5 cm soil depth than the 5-15, 15-30 and 30-100 cm depths and all other soils (Table 2). Soil was greater in the in the 0-5 cm, 5-15 cm and 15-30 cm depths in the IP than the IMP or ICT treatments.

Organic C contained in above-ground vegetation was greater on NSB sites than IP; however, IP biomass was removed by grazing. Crops were not considered as permanent vegetation. Prior to adjustment for agricultural CO₂ emissions, (total), soil C and C on site was greatest to least in the order IP>ICT>IMP>NSB (Table 3). Conversion of land to IP resulted in less C emitted to the atmosphere than IMP- or ICT- managed crops because less fertilizer and farm operations were necessary. After adjustment for agricultural CO₂ emissions, (net) C in soils was greatest to least in the order IP>ICT>NSB>IMP.

We estimated that if NSB sites were converted to IMP a net loss of 0.15 kg C m⁻² over 30 years would occur (Table 3). We estimated a net gain of 0.80 kg C m⁻² over 30 years if NSB sites were converted to ICT, and if converted to IP one could expect net gain of 3.56 kg C m⁻² over a 30 year period. We estimated that if IMP was converted to ICT a net gain of 0.95 kg C m⁻² over 30 years would occur. If IMP land was converted to IP an estimated net gain of 3.71 kg C m⁻² over a 30 year period would occur.

Manag

**Table 2. Ba
native sage**

Tre:

Native sage

Irrigated mc
plow crop

Irrigated cor
tilled crop

Irrigated pas

† In each column,
Square Mean
‡ Statistical comp
significant at

Regional and

Changes in agri
selected land n
impact to landc
managed proper
pollution. Estim
because the dyn
understood. Sto
ecosystems beca
values obtained
globally in soils
NSB, 2) all land
was converted t
increasing popul
converted to IP

Table 2. Bulk density, Walkley Black C and loss on ignition C (LAI-C) in soils growing native sagebrush, irrigated moldboard plowed cropland, irrigated conservation tilled cropland and irrigated pastures in Southern Idaho.

Treatment	Soil Depth	Bulk Density	WBC _†	LAI-C
		-- Mg m ⁻³ --	----- g C kg ⁻¹ soil-----	
Native sagebrush	0-5	0.97 b	12.7 a	10.1 a
	5-15	1.28 ab	4.7 c	4.1 c
	15-30	1.34 a	5.4 c	5.1 c
	30-100	1.36 a	4.0 c	3.5 c
Irrigated moldboard plow crop	0-30 ‡	1.28 ab	7.8 b	6.9 b
	30-100	1.37 a	6.0 bc	5.1 bc
Irrigated conservation tilled crop	0-15 ‡	1.38 a	8.9 b	7.6 b
	15-30 ‡	1.38 a	6.9 bc	6.8 bc
	30-100	1.37 a	3.7 c	3.1 c
Irrigated pasture	0-30 ‡	1.33 a	8.5 b	7.9 b
	30-100	1.40 a	4.3 c	3.8 c

† In each column, values followed by the same letter are not significantly different as determined by the Least Square Means Test ($P \leq 0.05$), $n = 30$.

‡ Statistical comparisons in the ANOVA showed that soil bulk density with respect to soil depth were not significant at $P \leq 0.05$. Therefore, data were combined (Snedecor and Cochran 1980; Kirk 1982).

Regional and Global Implications

Changes in agricultural practices have great potential to sequester C. In most cases converting selected land managed as IMP to ICT or IP can be implemented with modest economic impact to landowners and pose relatively few socioeconomic issues. If agricultural land is managed properly, these practice shifts would also potentially reduce erosion and water or air pollution. Estimating the potential for C sequestration in terrestrial ecosystems is difficult because the dynamics that control C flow among plants, soils and the atmosphere are poorly understood. Storage of C in below ground systems is the best long-term option in terrestrial ecosystems because C in soils has a longer residence time than most plant biomass. Using the values obtained in southern Idaho, we estimated C storage in soils locally, regionally and globally in soils if: 1) 10% of irrigated land now in IMP agriculture was converted back to NSB, 2) all land presently in IMP was converted to ICT and 3) 10 % of land in irrigated IMP was converted to IP. Since increased agricultural production will be necessary to feed an increasing population, it is impractical to suggest that a large portion of land in IMP can be converted to IP

to account for a
ed in the Pacific
l worldwide was
and in each area.
st, 24,322,029 ha
e, 1994; Howell,
projected to be
iding the Mg C
ease of CO₂ - C

(ANOVA) for a
Residuals were
programs (SAS
nce of treatment

, site C, net C in
Therefore, results
980; Kirk 1982).
density was less
and all other soils
5-30 and 30-100
5-15 cm and 15-

B sites than IP;
ed as permanent
C and C on site
on of land to IP
ops because less
gricultural CO₂
P.

kg C m⁻² over 30
30 years If NSB
of 3.56 kg C m⁻²
t gain of 0.95 kg
ated net gain of

The reported amounts of C stored in native sagebrush vegetation and irrigated agricultural systems are similar throughout the United States and worldwide (Bowman et al., 1999; Collins et al., 1999; Amthor and Huston, 1998; Potter et al., 1998; Rasmussen and Parton, 1994; Schlesinger, 1977;). These data were used to calculate potential C storage for irrigated agriculture in the Pacific Northwestern U.S., the Western U.S., and worldwide over a 30 year period. If land currently in IMP is converted to NSB we estimated a gain of 1.5 Mg C ha⁻¹ (Table 4). Little of this land is managed with conservation tillage. We estimate an increase of 9.5 Mg C ha⁻¹ over 30 years if the land presently managed with IMP were converted to ICT. Using this value we calculated that 8.6×10^7 Mg C (0.15% of the total C emitted in the next 30 yr) could potentially be sequestered in irrigated soils in the Pacific Northwestern U.S (Table 4). Using these values to represent C gains for all irrigated crop land in the western U.S. and if land in IMP were converted to ICT, a possible 2.3×10^8 Mg C (0.40% of the total C emitted in the next 30 yr) could be sequestered in irrigated agricultural soils in the next 30 years. If the world's IMP land were converted to ICT, 2.5×10^9 Mg C (4.38% of the total C emitted in the next 30 yr) could be sequestered in the next 30 years. A shift of 10 % of current IMP land to ICT is a reasonable conservation practice goal. Similarly, we can not expect all land presently in IMP to be converted to IP, but a 10 % conversion is feasible. If we predict a storage increase of 37.1 Mg C ha⁻¹ for IMP converted to IP and assume 10 % of IMP land is converted to IP, an estimated 3.4×10^7 Mg C (0.05% of the total C emitted in the next 30 yr) could be sequestered in Pacific Northwestern U.S. soils and 9.0×10^7 Mg C (0.16% of the total C emitted in the next 30 yr) in the Western U.S. soils in the next 30 years. If our study values are used to represent C gains for irrigated crop land worldwide, an estimated 9.6×10^8 Mg C (1.68% of the total C emitted in the next 30 yr) could be sequestered if irrigated land presently managed as IMP were converted to ICT (Table 4). Conversion of crop land to irrigated pasture could also relieve grazing pressure on public rangelands, an issue of heated debate between environmentalists and ranchers.

If irrigated agriculture is expanded due to increase in water use efficiency, one could expect a gain in C sequestration. If NSB were converted to ICT, 8.0 g C ha⁻¹ could be sequestered (Table 4). Predicting a storage increase of 8.0 Mg C ha⁻¹ for NSB converted to ICT and assuming 10 % expansion of irrigated agriculture, an estimated 7.2×10^6 Mg C (0.01% of the total C emitted in the next 30 yr) could be sequestered in Pacific Northwestern U.S. soils and 1.9×10^7 Mg C (0.033% of the total C emitted in the next 30 yr) in the Western U.S. soils in the next 30 years. If our study values are used to represent C gains for irrigated crop land worldwide, an estimated 2.1×10^8 Mg C (0.37% of the total C emitted in the next 30 yr) could be sequestered (Table 4). If NSB were converted to IP, 35.6 g C ha⁻¹ could be sequestered. Predicting a storage increase of 35.6 Mg C ha⁻¹ for NSB converted to IP and assuming 10% expansion of irrigated agriculture, an estimated 3.2×10^8 Mg C (0.5% of the total C emitted in the next 30 yr) could be sequestered in Pacific Northwestern U.S. soils and 8.7×10^8 Mg C (1.53% of the total C emitted in the next 30 yr) in the Western U.S. soils in the next 30 years. If our study values are used to represent C gains for irrigated crop land worldwide, an estimated 9.3×10^9 Mg C (16.3% of the total C emitted in the next 30 yr) could be sequestered

Table 3. Organic carbon in: soils¹, above ground biomass² and on sites at present³, carbon emitted during agricultural operations⁴, net organic carbon in soils⁵, and net carbon gain on sites⁶. †

Vegetation	Carbon Present	Carbon Emitted	Net Carbon Gain
	Soil †	Soil †	Soil †
	Above	Above	Above

on and irrigated
e (Bowman et al.,
; Rasmussen and
tial C storage for
d worldwide over
a gain of 1.5 Mg
. We estimate an
l with IMP were
5% of the total C
oils in the Pacific
irrigated crop land
: 2.3 x 10⁸ Mg C
gated agricultural
T, 2.5x10⁹ Mg C
: next 30 years. A
ce goal. Similarly,
0% conversion is
nverted to IP and
0.05% of the total
m U.S. soils and
n U.S. soils in the
irrigated crop land
next 30 yr) could
to ICT (Table 4).
ressure on public
rs.
ciency, one could
; C ha⁻¹ could be
NSB converted to
ed 7.2x10⁶ Mg C
ific Northwestern
yr) in the Western
gains for irrigated
tted in the next 30
g C ha⁻¹ could be
nverted to IP and
lg C (0.5% of the
tern U.S. soils and
m U.S. soils in the
irrigated crop land
: next 30 yr) could

Table 3. Organic carbon in: soils¹, above ground biomass² and on sites at present³, carbon emitted during agricultural operations⁴, net organic carbon in soils⁵ and net carbon gain on sites⁶. †

Vegetation	Carbon Present			Net Carbon Gain		
	Soil ¹ ‡	Above ground ² ¶	Site ³	Carbon emitted ⁴ §	Soil ⁵ ‡	Site ⁶
kg C m ⁻²						
Native sagebrush	5.91 c	0.42 a	6.34 c	0.00 d	5.91 c	6.34 c
Irrigated moldboard plow crops	7.29 b	0.00 c	7.29 b	1.10 a	6.19 b	6.19 c
Irrigated conservation till crops	8.01 b	0.00 c	8.01 b	0.87 b	7.14 b	7.14 b
Irrigated pasture	10.14 a	0.05 b	10.19 a	0.29 b	9.85 a	9.90 a

† In each column, values followed by the same letter are not significantly different as determined by the Least Square Means Test (*P* # 0.05), *n* = 30.

‡ Values of organic C stored in soils are based on the Walkley Black procedure.

¶ Carbon in soils, above ground vegetation and on the sites at the present time.

§ Estimated carbon emitted in production of fertilizer, fuel consumption in farm operations and via irrigation water over a 30-year period.

Table 4. Potential organic carbon gain by conversion of irrigated lands currently in moldboard plowing systems to conservation tillage, conversion of native sagebrush to irrigated conservation tillage, conversion of native sagebrush to irrigated pasture and conversion of 10% of irrigated lands currently in moldboard plowing systems to irrigated pasture over the next 30 years.

Vegetation Conversion	C gained from a 10 % conversion		Pacific Northwest United States †		Western United States ‡		Worldwide †		
	-Mg C ha ⁻¹ -	Mg C	% C _s /C _{EW} §	Mg C	% C _s /C _{EW} §	Mg C	% C _s /C _{EW} §	Mg C	% C _s /C _{EW} §
Irrigated moldboard plow to irrigated conservation tillage †	9.5	8.6 x 10 ⁷	0.15	2.3x10 ⁸	0.40	2.5x10 ⁹	4.38		
Native sagebrush to irrigated conservation tillage	8.0	7.2x10 ⁶	0.01	1.9x10 ⁷	0.03	2.1x10 ⁸	0.37		
Native sagebrush to irrigated pasture	3.56	3.2x10 ⁸	0.56	8.7x10 ⁸	1.53	9.3x10 ⁹	16.32		
10% of irrigated moldboard plow to irrigated pasture †	37.1	3.4 x 10 ⁷	0.06	9.0 x 10 ⁷	0.16	9.6 x 10 ⁸	1.68		

† Estimated C gain from 100% conversion of moldboard plow to conservation tillage and 10% conversion of moldboard plow agriculture to irrigated pasture.

‡ Land area in irrigated cropland in Pacific Northwest - 9, 055,979 ha, Western United States 24,322,029 ha, worldwide 260,000,000 ha.

§ %C_s/C_{EW} = carbon sequestered (C_s) divided by the amount of C projected to be emitted worldwide during the next 30 yr, which is 5.7 x 10¹⁰ Mg C (C_{EW}) multiplied by 100.

Table 5. Potential carbon transfer by converting an equal amount (10%) of rainfed moldboard plow land back to native forest or grassland on a basis of 1 unit of irrigated rainfed agricultural land to 1 unit of native forest or grassland and conversion of equal amount (10%) of rainfed moldboard plow land back to native forest or grassland on the basis of 1 unit of irrigated rainfed agricultural land to 2 units of native forest or grassland

Table 5. Potential carbon transfer by converting an equal amount (10%) of rainfed moldboard plow land back to native forest or grassland on a basis of 1 unit of irrigated rainfed agricultural land to 1 unit of native forest or grassland and conversion of equal amount (10%) of rainfed moldboard plow land back to native forest or grassland on the basis of 1 unit of irrigated rainfed agricultural land to 2 units of native forest or grassland

Conversion of Vegetation	C Stored From Conversion		Pacific Northwest United States		Western United States		Worldwide	
	Mg C ha ⁻¹	Mg C	%C _s /C _{EW} †	Mg C	%C _s /C _{EW} †	Mg C	%C _s /C _{EW} †	
Rainfed moldboard plow to native forest on a 1 unit:1 unit basis	5.6	5.1 x 10 ⁷	0.09	1.4 x 10 ⁸	0.24	1.5 x 10 ⁹	2.63	
Rainfed moldboard plow to native grassland on a 1 unit:1 unit basis	13.0	1.2 x 10 ⁸	0.21	3.2 x 10 ⁸	0.56	3.4 x 10 ⁹	5.96	
Rainfed moldboard plow to native forest on a 2 unit:1 unit basis	5.6	1.1 x 10 ⁸	0.18	2.8 x 10 ⁸	0.49	3.0 x 10 ⁹	5.26	
Rainfed moldboard plow to native grassland on a 2 unit:1 unit basis	13.0	2.4 x 10 ⁸	0.42	6.4 x 10 ⁸	1.20	6.8 x 10 ⁹	11.93	

† %C_s/C_{EW} = carbon sequestered (C_s) divided by the amount of C projected to be emitted worldwide during the next 30 yr, which is 5.7 x 10¹⁰ Mg C (C_{EW}) multiplied by 100.

Since the earth releases 1.9×10^9 Mg C yr⁻¹ (Schlesinger, 1995; Amthor and Huston, 1998), the conversion of 10% IMP to IP, resulting in a possible 9.6×10^8 Mg C sequestered over 30 yrs, may be insignificant (Table 4). However, if crops were produced via high output irrigated agriculture while less productive rainfed agricultural land were returned to temperate forest or native grassland, an increase of 5.6 and 13 Mg C ha⁻¹, respectively, could be gained over 30 years for each unit of rainfed land converted to native vegetation (Table 5). The amount of irrigated agriculture can likely be increased at least 10% solely through increases in irrigation efficiency and waste water reuse (Howell, 2000). Using a conversion basis of 1 unit of irrigated agriculture to return 1 unit of rainfed agricultural land to native forest, if irrigated agriculture were expanded 10% (meaning that an additional 2.6×10^7 ha of arid or semiarid land were irrigated) and the equal amount of land being managed as rainfed agricultural land were converted to native forest, there is potential to sequester 5.1×10^7 Mg C (0.09% of the total C emitted in the next 30 yr) in the Pacific North West, 1.41×10^8 Mg C (0.24% of the total C emitted in the next 30 yr) in the western U.S. and 1.5×10^9 Mg C (2.6% of the total C emitted in the next 30 yr) worldwide (Table 5). If the rainfed agricultural land were converted to native grassland, there is a potential to sequester 1.2×10^8 Mg C (0.2% of the total C emitted in the next 30 yr) in the PNW, 3.2×10^8 Mg C (0.6% of the total C emitted in the next 30 yr) in the western U.S. and 3.4×10^9 Mg C (5.9% of the total C emitted in the next 30 yr) worldwide (Table 5).

However, irrigated agricultural land typically produces twice the crop yield of rainfed agricultural land (Bucks et al., 1990; Howell, 2000). If irrigated agriculture were expanded 10%, each hectare of new irrigated land could produce the same crop yield as 2 ha of rainfed land (Bucks et al., 1990; Tribe, 1994; Howell, 2000). Under this scenario, the conversion of irrigated land to native forest could potentially sequester 1.0×10^8 Mg C (0.2% of the total C emitted in the next 30 yr) in the PNW, 2.8×10^8 Mg C (0.5% of the total C emitted in the next 30 yr) in the western U.S. and 3.0×10^9 Mg C (5.26% of the total C emitted in the next 30 yr) worldwide (Table 5). If converted to native grassland in this 2:1 conversion scenario, there is a potential to sequester 2.4×10^8 Mg C (0.4% of the total C emitted in the next 30 yr) in the PNW, 6.4×10^8 Mg C (1.2% of the total C emitted in the next 30 yr) in the western U.S. and 6.8×10^9 Mg C (11.9% of the total C emitted in the next 30 yr) worldwide. If highly erosive rainfed lands were selected or if rainfed lands urgently needed for habitat restoration were targeted for such a conversion, significant additional erosion, water quality and habitat benefits could also result.

Since native desert or semi-desert has relatively little ecosystem C compared to forest, grassland or wetland ecosystems (Houghton et al., 1999; Amthor and Huston, 1998; Schlesinger, 1977), converting irrigated moldboard plow crop land back to desert or semi-desert could result in a soil C gain of 0.15 kg C m^{-2} . A sequestration of only 1.90×10^6 Mg C 30 yr⁻¹ might be expected in the Pacific Northwestern U.S., 5.10×10^6 Mg C 30 yr⁻¹ in the western U.S. and 5.46×10^7 Mg C 30 yr⁻¹ worldwide. This modest C accumulation in soil would be tempered by the fact that the conversion would require substantial policy incentives and decades to implement. Substantially more C may be sequestered by selectively returning rainfed agricultural land derived from forest, grassland or wetlands back to native vegetation. Tropical and temperate forests typically contain from 10-12 kg C m⁻², grasslands contain from 18-20 kg C m⁻², and wetland ecosystems contain from 60-70 kg C m⁻², whereas arid and semi-arid lands contain 5-7 kg C m⁻² (Houghton et al., 1999; Ross et al., 1999; Schimel et al., 2000). Since nearly a third of the yield and nearly half of the value of crops in the U.S. are

produced on irrigated land (Tribe, 1990; Tribe, 1994), the conversion of irrigated agriculture to native vegetation, accompanied by the restoration of wetlands back to

Factors Affecting

Grazing affects the distribution of C in ecosystems (Williams, 1990). Williams (1990) found that grazing with herbivores had little effect on C sequestration, but it was a part of long-term

We recognize the need for more precise estimates of C sequestration on a worldwide basis. There are many different scenarios with potential for the gain of C. Improving irrigation management that reduces erosion and world-wide uses of C via erosion

Flood and microorganisms (Sojka et al., 1999) transport off site leaching (Aase et al., 1999) systems to reduce polyacrylamide, (1998). Additional technology needed in agriculture in the to erosion. Because occurred in recent conservative.

Conclusions

As ecosystems are controlled by cli

1991; Harmon et al., 1990). Therefore, at equilibrium, the amount of C added to the soil via vegetation is equal to the amount of C lost through organic matter degradation and other losses (Henderson, 1995) and eventually a maximum limit will be reached. Although irrigated agricultural systems can not accrue C indefinitely, with improved management they can potentially remove substantial amounts of C from the atmosphere for the next 30-50 years. The potential C sequestered on site by conversion of native vegetation to irrigated agriculture is above the steady state equilibrium of native vegetation. This is in contrast to rainfed agricultural systems which are currently attempting modest C gains to attain near-baseline C concentrations by implementing reduced or no-tillage practices. Rainfed agricultural lands with or without no-till practices have soil C values far below those of native vegetation. A third of the yield and nearly half of the value of crops in the U.S. and worldwide are produced on the irrigated 15-17% of arable lands that are largely in arid or semi-arid climatic zones (Tribe, 1994; Bucks et al., 1990). Since the earth releases 1.9×10^9 Mg C yr⁻¹ (Amthor and Huston, 1998; Schlesinger, 1995), the conversion of 10 % IMP to ICT resulting in a possible 2.5×10^9 Mg C sequestered over 30 years is a modest 4.47% compared to the total C released into the atmosphere over that time period. However, if crops were produced via high output irrigated agriculture, while selected less-productive rainfed agricultural land were returned to temperate forest or native grassland on a world-wide basis, there could be substantial reductions in atmospheric CO₂. Policy makers and agricultural research infrastructure should recognize the enormous potential benefit of land and water management strategies, policies and incentives that could expand arid zone irrigated agriculture as a means for efficient food and fiber production along with substantial C sequestration potential. This potential would be enhanced if coupled with selective return of less efficient rainfed agricultural lands derived from forest, grassland or wetlands back to native vegetation. We recognize that such an expansion would have to be accompanied by renewed efforts of water development. While water resource development has been occurring at a modest pace worldwide since 1990, Howell (2000) indicated the potential for increased extent of irrigation via efficiency improvements and waste water use. Recognition of these potential C benefits should provide an incentive to fund research and pursue management strategies that are possible without sacrificing production and which could increase restoration of native ecosystems, reduce erosion and improve water quality through appropriate targeting of the strategy.

References

- Aase, J. K., D. L. Bjorneberg, and R.E. Sojka, 1998. Sprinkler irrigation runoff and erosion control with polyacrylamide - laboratory tests. *Soil Sci. Soc. Am. J.* **62**:1681-1687.
- Amthor, J.S. and M.A. Huston 1998. Terrestrial ecosystem responses to global change: a research strategy, ORNL/TM-1998/27, Oak Ridge National Laboratory. 157 pp.
- Bailey, R.G. 1998. Ecoregions of North America. U.S. Department of Agriculture. Forest Service, US Government Printing Office, Washington D.C.
- Birch, H.F. and M.T. Friend. 1956. The organic matter and nitrogen status of east Africa soils. *J. Soil. Sci.* **7**:156-167.
- Blake, G. R. and K.H. Hartage. 1982. Bulk density. In p. 363-375 A.L. Page, R.H. Miller, and D.R. Keeney. (eds). *Methods of soil analysis. Part 2. Chemical and microbiological properties.* *Am. Soc. Agron.* Madison, WI.

- Bowman, R.A.,
changes in i
- Bruce, J.P., M.
sequestratio
- Bucks, D.A., T.
Managemen
(eds). *Am. So*
- Burke, J.C., C.
Texture, cli
soils. *Soil Sc*
- Cambardella, C.
grassland cu
- Collett, R.A. 199
Resources (C
182 pp.
- Collins, H.P., R.
E.A. Paul. 1
J. **63**:584-59
- Dewar, R.C., 19
of managed
- Dommar, J.F., a
properties of
- Entry, J. A., and
remove mic
1905-1914.
- Entry, J. A., an
Douglas-fir
- Follett, R.F. 200
Tillage & Re
- Frank, A.B., D.I
Northern Gr
48:470-474.
- Harmon, M.E., V
old growth f
- Henderson, G.S
productivity
functions in
- Hontoria, C., J.C
carbon and s
- Howell, T.A. 20
B.L. Benhar
of Engineers
- Houghton, R.A
contribution.

ed to the soil via
dation and other
Although irrigated
gement they can
next 30-50 years.
irrigated agriculture
ntrast to rainfed
near- baseline C
gricultural lands
ive vegetation. A
vide are produced
id climatic zones
yr⁻¹ (Amthor and
ting in a possible
e total C released
d via high output
were returned to
ld be substantial
astructure should
strategies, polices
for efficient food
otential would be
ral lands derived
ize that such an
velopment. While
wide since 1990,
n via efficiency
ts should provide
possible without
osystems, reduce
gy.

unoff and erosion
681-1687.
global change: a
157 pp.
griculture. Forest
us of east Africa
s, R.H. Miller, and
d microbiological

- Bowman, R.A., M.F. Vigil, D.C. Nielsen, and R.L. Anderson. 1999. Soil organic matter changes in intensively cropped dryland systems. *Soil Sci. Soc. Am. J.* **63**:186-191.
- Bruce, J.P., M. Frome, E. Haites, H. Janzen, R. Lal and K. Paustian. 1999. Carbon sequestration in soils. *J. Soil Water Conserv.* **59**:382-389.
- Bucks, D.A., T.W. Sammis, and G.L. Dickey. 1990. Irrigation for arid areas. pp 449-548. In: Management of Farm Irrigation Systems. G.J. Hoffman, T.A. Howell and K.H. Solomon (eds). *Am. Soc. Ag. Eng.*, St. Joseph, MI.
- Burke, J.C., C.M. Yonker, W.J. Parton, C.V. Cole, K. Flach, and D.S. Schimel. 1989. Texture, climate and cultivation effects on soils organic matter content in U.S. grassland soils. *Soil Sci. Soc. Am. J.* **53**:800-805.
- Cambardella, C.A., and E.T. Elliott. 1992. Particulate soil organic matter changes across a grassland cultivation sequence. *Soil Sci. Soc. Am. J.* **56**:777-783.
- Collett, R.A. 1982. Soil Survey of Ada County area. U.S. Department of Agriculture, Natural Resources Conservation Service., US Government Printing Office, Washington D.C. 182 pp.
- Collins, H.P., R.L. Blevins, L.G. Bundy, D.R. Christenson, W.A. Dick, D.R. Huggins and E.A. Paul. 1999. Soil carbon dynamics in corn-based agroecosystems: *Soil Sci. Soc. Am. J.* **63**:584-591.
- Dewar, R.C., 1991. Analytical model of carbon storage in the trees, soils, and wood products of managed forests. *Tree Physiol.* **8**: 239-258.
- Dommar, J.F., and W.D. Williams. 1990. Effect of grazing and cultivation on some chemical properties of soils in the mixed prairie. *J. Range. Manage.* **43**:456-460.
- Entry, J. A., and R. E. Sojka, 2000. The efficacy of polyacrylamide and related compounds to remove microorganisms and nutrients from animal wastewater. *J. Environ. Qual.* **29**: 1905-1914.
- Entry, J. A., and W.H. Emmingham. 1998. Influence of forest age on forms of carbon in Douglas-fir soils in the Oregon Coast Range. *Can. J. For. Res.* **28**: 390-395.
- Follett, R.F. 2001. Soil management concepts and carbon sequestration in cropland soils. *Soil Tillage & Research.* **61**:77-92.
- Frank, A.B., D.L. Tanaka, L. Hofmann and R.F. Follett. 1995. Soil carbon and nitrogen of Northern Great Plains grasslands as influenced by long term grazing. *J. Range. Manage.* **48**:470-474.
- Harmon, M.E., W. Ferrell and J.F. Franklin. 1990. Effects on carbon storage of conversion of old growth forests to young growth forests. *Science* **247**: 699-702.
- Henderson, G.S. 1995. Soil organic matter: a link between forest management and productivity. In p 419-436. W.F. McFee and J. M. Kelley (eds.) Carbon forms and functions in soils. Soil Science Society of America Press., Madison, WI.
- Hontoria, C., J.C. Rodriguez-Murillo and A. Saa. 1999. Relationships between soil organic carbon and site characteristics in peninsular Spain. *Soil Sci. Soc. Am. J.* **63**:614-621.
- Howell, T.A. 2000. Irrigations role in enhancing water use efficiency. In: p 67-80. R.G. Evans, B.L. Benham and T.P. Trooien (eds). National Irrigation Symposium. American Society of Engineers, St. Joseph, MI.
- Houghton, R.A., J.L. Hackler and K.T. Lawrence. 1999. The U.S. carbon budget: contributions from land use change. *Science* **285**:574-578.

- Janzen, H.H., C.A. Campbell, E.G. Gregorich and B.H. Ellert. 1997. Soil carbon dynamics in Canadian ecosystems. p 57-80. In: R. Lal, J. Kimble, E. Levine and B.A. Stewart (eds.) *Soils and global change*. CRC Press, Boca Raton, FL.
- Jobbagy, E.G., and R.B. Jackson. 2000. The vertical distribution of organic carbon and its relation to climate and vegetation. *Ecol. Appl.* **10**: 423-436.
- Johnson, D. W. 1992. Effects of forest management on soil carbon storage. *Water, Air, Soil Pollut.* **64**: 83-120.
- Kimble, L., and B.A. Stewart. 1995. World soils as a source or sink for radiatively- active gasses. In: p 1-7 R. Lal, J. Kimble, E. Levine and B.A. Stewart R. (eds.) *Soils and global change*. CRC Press, Boca Raton, FL.
- Kirk, R.E., 1982. *Experimental Design: Procedures for the Behavioral Sciences*. 2nd ed. Brooks/Cole Publishing, Monterey, CA, 911 pp.
- Lal, R., R.F. Follett, J. Kimble and C.V. Cole. 1999. Managing U.S. cropland to sequester carbon in soil. *J. Soil Water Conserv.* **59**:374-381.
- Liski, J., H. Iivesniemi, A. Makela and C.J. Westman. 1999. CO₂ emissions from soil in response to climatic warming are overestimated - the decomposition of old soil organic matter is tolerant of temperature. *Ambio* **28**:171-174.
- Nelson, D.W., and L.E. Sommers. 1996. Total carbon, organic carbon and organic matter. In p 961-1010. *Methods of soil analysis. Part 3, Chemical and microbiological properties*. J.M. Bigham (ed.). ASA, CSSA, SAAJ, Madison, WI.
- Paustian, K., O. Anderon, H. Janzen, R. Lal, P. Smith, G. Tian, H. Tiessen, M. van Noordwijk and P. Woormer. 1997. Agricultural soil as a C sink to offset CO₂ emissions. *Soil Use and Management* **13**:230-244.
- Potter, K.N., H.A. Torbert, O.R. Jones, J.E. Matocha, J.E. Morrison, Jr., and P.W. Unger. 1998. Distribution and amount of soil organic C in long term management systems in Texas. *Soil & Tillage Research.* **47**: 309-321.
- Rasmussen, P.E., and H.P. Collins. 1991. Long-term impacts of tillage, fertilizer and crop residue on soil organic matter in temperate semi-arid regions. *Adv Agron.* **45**:93-134.
- Rasmussen, P.E., and W.J. Parton. 1994. Long-term effects of residue management in wheat-fallow: I. Inputs, yield, and soil organic matter. *Soil Sci. Soc. Am. J.* **58**: 523-530.
- Ross, D.J., K.R. Tate, N.A. Scott and C.W. Feltham. 1999. Land-use change: effects on soil carbon, nitrogen and phosphorus pools and fluxes in three adjacent ecosystems. *Soil Biology and Biochemistry* **31**: 803-813.
- SAS Institute Inc. 1996. *SAS User's Guide: Statistics - Version 6.03 Edition*. Statistical Analysis System (SAS) Institute Inc., Cary, NC. 584 pp.
- Schimel, D., J. Melillo, H. Tian, A.D. McGuire, D. Kicklighter, T. Kittel, N. Rosenbloom, S. Running, P. Thornton, D. Ojima, W. Parton, R. Kelly, M. Sykes, R. Neilson and B. Rizzo. 2000. Contribution of increasing CO₂ and climate to carbon storage by ecosystems in the United States. *Science* **287**:2004-2006.
- Schlesinger, W.M. 1977. Carbon balance in terrestrial detritus. *Ann. Rev. Ecol.* **8**:51-81.
- Schlesinger, W.M. 1990. Evidence from chronosequence studies for a low carbon-storage potential of soils. *Nature* **348**:232-234.
- Schlesinger, W.M. 1995. An overview of the C cycle. p 9-26. In R. Lal, J. Kimble, E. Levine and B.A. Stewart. (eds.) *Soils and global change*. CRC Press, Boca Raton, FL
- Schlesinger, W.M. 1999. Carbon sequestration in soils. *Science* **284**:2095.

- Schuman, G.E., grazing ma
Ecological
- Snedecor, W.G.
Press, Ames
- Sojka, R.E., R.
Polyacrylan
54:325-331
- Sojka, R.E., R.I
multiple ap
62:1672-16
- Sojka, R. E. a
movement c
- Tribe, D. 1994.
Internationa
- Van Cleve, K.
developmen
941-955.
- Wang, Y., R. A
turnover in :
- West, T.O. and
net carbon
Ecosyst. En

arbon dynamics in
A. Stewart (eds.)

ic carbon and its

e. *Water, Air, Soil*

radiatively- active

) *Soils and global*

Sciences. 2nd ed.

pland to sequester

sions from soil in

of old soil organic

organic matter. *In*

ological properties.

Tiessen, M. van

set CO₂ emissions.

and P.W. Unger.

gement systems in

fertilizer and crop

m. **45**:93-134.

agement in wheat-

523-530.

age: effects on soil

t ecosystems. *Soil*

Edition. Statistical

N. Rosenbloom, S.

R. Neilson and B.

arbon storage by

col. **8**:51-81.

ow carbon-storage

Kimble, E. Levine

on, FL

Schuman, G.E., J.D. Reeder, J.T. Manley, R.J. Hart, and W.A. Manley. 1999. Impact of grazing management on the carbon and nitrogen balance of a mixed-grass rangeland. *Ecological Applications*. **9**:65-71.

Snedecor, W.G. and W.G. Cochran. 1980. *Statistical methods*. 7th ed. Iowa State University Press, Ames, Iowa. 354 p.

Sojka, R.E., R.D. Lentz, C.W. Ross, T.J. Trout, D.L. Bjorneberg, and J.K. Aase. 1998a. Polyacrylamide effects on infiltration in irrigated agriculture. *J. Soil. Water. Cons.* **54**:325-331.

Sojka, R.E., R.D. Lentz, and D.T. Westermann, 1998b. Water and erosion management with multiple applications of polyacrylamide in furrow irrigation. *Soil Sci. Soc. Am. J.* **62**:1672-1680.

Sojka, R. E. and J.A. Entry. 2000. Influence of polyacrylamide application to soil on movement of microorganisms in runoff water. *Environ. Pollut.* **108**:405-412.

Tribe, D. 1994. Feeding and Greening the World, the Role of Agricultural Research. CAB International. Wallingford, United Kingdom. 274pp.

Van Cleve, K., C.T. Dryness, G.M. Marion, and R. Erickson. 1993. Control of soil development on the Tanana River floodplain, interior, *Alaska.Can. J. For. Res.* **23**: 941-955.

Wang, Y., R. Amundson and S. Trumbore. 1999. The impact of land use change on C turnover in soils. *Gobal Biogeochemical Cycles*.**13**:47-57.

West, T.O. and G. Marland. 2001. A synthesis of carbon sequestration, carbon emissions, and net carbon flux in agriculture: comparing tillage practices in the United States. *Agric. Ecosyst. Environ.* **91**: 217-232.